

Preoperative Patient Orders
Pre-Printed Physician's Orders

Patient Name: _____

Surgeon: _____

PRE OPERATIVE MEDICATIONS

- | | |
|--|--|
| <input type="checkbox"/> acetaminophen 1000 mg PO
<input type="checkbox"/> gabapentin 300 mg PO
<input type="checkbox"/> celecoxib (Celebrex) 400 mg PO
Other: _____
_____ | <input type="checkbox"/> metronidazole (Flagyl) 1 g PO 30 min preop
<input type="checkbox"/> phenazopyridine (Pyridium) 100 mg PO 30-60 min preop
<input type="checkbox"/> ciprofloxacin 500 mg PO 1-2 h preop
<input type="checkbox"/> misoprostol (Cytotec) 400 mcg buccally 2 h preop
<input type="checkbox"/> doxycycline 100 mg PO 30 min preop |
|--|--|

INTRA OPERATIVE MEDICATIONS

If NO allergy or non-life threatening penicillin/beta-lactam allergy

- cefazolin (Ancef) 2 g IV (if weight less than 120 kg)
- cefazolin (Ancef) 3 g IV (if weight 120 kg or greater)
- metronidazole (Flagyl) 500 mg IV

If anaphylaxis to penicillin/beta-lactams

- vancomycin 1 g IV (if weight 80 kg or less). Begin infusion 60 min prior to time of surgery.
- vancomycin 1.25 g IV (if weight 80.1 to 99.9 kg). Begin infusion 90 min prior to time of surgery.
- vancomycin 1.5 g IV (if weight 100 kg or greater). Begin infusion 90 min prior to time of surgery.
- Vancomycin patients :** Initiate IV LR at 100 mL/h with # 20 gauge or greater in left hand/arm(unless operative limb)
- clindamycin 600 mg IV
- ciprofloxacin 400 mg IV
- Other _____
- gentamicin _____ (2 mg/kg) IV
- ceftriaxone 2 g IV

ADDITIONAL TESTING (ABOVE MEDICAL DIRECTIVE)

FASTING INSTRUCTIONS

- | | |
|---|--|
| <input type="checkbox"/> Routine diet OR <input type="checkbox"/> Clear fluids day before surgery
<input type="checkbox"/> Clear fluids for 24 to 48 hours before surgery
<input type="checkbox"/> NPO after midnight (may have clear fluids until 4 hours before surgery) | ERAS <input type="checkbox"/> up to 800 mL carbohydrate rich drink HS
<input type="checkbox"/> Clear fluids until 4 hours before surgery
<input type="checkbox"/> 400 mL carbohydrate rich drink 4 hours preop |
|---|--|

INTERVENTIONS

- Clip hair from operative site
- apply bilateral knee high stockings prior to going into OR
- apply thigh high SCD sleeves in OR (Day Surgery to measure preoperatively)
- Other _____

BOWEL PREPARATION

- | | |
|--|---|
| <input type="checkbox"/> No bowel prep
<input type="checkbox"/> Lyte prep 4L PO (polyethylene glycol/electrolyte solution) as per instructions
<input type="checkbox"/> sodium phosphate enema (Fleet enema) PR X 1 in the morning on the day of surgery | <input type="checkbox"/> sodium picosulfate (Pico-Salax) in 150mL water, PO X 2 doses, first sachet given at 1200 and second given at 1700, both on the day before surgery
<input type="checkbox"/> bisacodyl (Dulcolax) 10 mg PO with each sachet of Pico-Salax |
|--|---|

ANTICOAGULATION

Instructions : _____

OR Consult Anesthesia **OR** Internal Medicine

CONSULTS

- | | | |
|--|-------------------------------------|---|
| <input type="checkbox"/> Internal Medicine | <input type="checkbox"/> Anesthesia | <input type="checkbox"/> Blood Conservation Nurse |
|--|-------------------------------------|---|

Date: _____ Time: _____ Physician Signature: _____

Physician Print Name: _____